



## Patterns and drivers of fish community assembly in a large marine ecosystem

Pécuchet, Lauréne; Törnroos, Anna; Lindegren, Martin

*Published in:*  
Marine Ecology - Progress Series

*Link to article, DOI:*  
[10.3354/meps11613](https://doi.org/10.3354/meps11613)

*Publication date:*  
2016

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Pécuchet, L., Törnroos, A., & Lindegren, M. (2016). Patterns and drivers of fish community assembly in a large marine ecosystem. *Marine Ecology - Progress Series*, 546, 239-248. <https://doi.org/10.3354/meps11613>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Patterns and drivers of fish community assembly in a large marine ecosystem

Laurène Pécuchet\*, Anna Törnroos, Martin Lindegren

\*Corresponding author: [laupe@aqua.dtu.dk](mailto:laupe@aqua.dtu.dk)

Marine Ecology Progress Series 546: 239–248 (2016)

Table S1. Species list and traits of demersal fish species present in the Baltic Sea International Trawl Survey (BITS) in Quarter 1 from 2003 to 2014. The ‘diet’ is taken from FishBase; ‘Lmean’ is the mean length of the species in the survey; ‘A50’ and ‘Fecundity’ are taken from Fishbase, ICES species sheet facts and stock assessment or from the literature as stated in Ref; ‘caudal’ and ‘body’ shape are derived from FishBase pictures.

Species	Diet	Lmean	A50	Fecundity	Caudal	Body	Ref	Area
<i>Agonus cataphractus</i>	benthivorous	135	2	3000	rounded	elongated	<sup>1</sup>	Baltic Sea
<i>Amblyraja radiata</i>	generalist	390	5.5	50	continuous	flat	<sup>2</sup>	North Sea
<i>Ammodytidae</i>	planktivorous	179	1.5	5000	forked	elongated	<sup>3</sup>	North Sea
<i>Anguilla anguilla</i>	generalist	570	12	2000000	continuous	eellike	<sup>4,5</sup>	Europe
<i>Arnoglossus laterna</i>	generalist	120	2	50000	rounded	flat	<sup>6</sup>	Scotland
<i>Buglossidium luteum</i>	benthivorous	100	3	13400	rounded	flat	<sup>7</sup>	English Channel
<i>Callionymus</i>	benthivorous	126	2.5	5000	rounded	normal	<sup>8</sup>	North Sea
<i>Chelidonichthys lucernus</i>	generalist	266	2	100000	truncated	normal	<sup>9</sup>	Mediterranean
<i>Cyclopterus lumpus</i>	planktivorous	205	5	100000	truncated	normal	<sup>10</sup>	Baltic Sea
<i>Enchelyopus cimbrius</i>	generalist	231	3	30000	rounded	elongated	<sup>11</sup>	?
<i>Eutrigla gurnardus</i>	generalist	170	3	14347	emarginated	normal	<sup>8</sup>	Irish Sea
<i>Gadus morhua</i>	piscivorous	303	3	1000000	truncated	normal	<sup>12</sup>	Baltic Sea
<i>Glyptocephalus cynoglossus</i>	benthivorous	230	4.5	100000	rounded	flat	<sup>13</sup>	Gulf of Maine
<i>Gobiidae</i>	generalist	73	1	4000*	rounded	normal	<sup>1</sup>	Baltic Sea
<i>Hippoglossoides platessoides</i>	generalist	160	10	380000	rounded	flat	<sup>14</sup>	Barents Sea
<i>Hyperoplus</i>	planktivorous	201	2	7000	forked	elongated	<sup>1</sup>	North Sea
<i>Lepidorhombus whiffiagonis</i>	generalist	129	2	300000	rounded	flat	<sup>15</sup>	Ireland
<i>Limanda limanda</i>	benthivorous	186	2.5	150000	rounded	flat	<sup>16</sup>	North Sea
<i>Lumpenus lampretaeformis</i>	benthivorous	226	3	1000	rounded	eellike	<sup>17</sup>	Barents Sea
<i>Melanogrammus aeglefinus</i>	generalist	222	2.5	535000	emarginated	normal	<sup>18,19</sup>	North Sea
<i>Merlangius merlangus</i>	piscivorous	208	2	400000	truncated	normal	<sup>20</sup>	Irish Sea
<i>Merluccius merluccius</i>	piscivorous	252	4.4	1000000	truncated	elongated	<sup>21,22</sup>	Bay of Biscay
<i>Microstomus kitt</i>	benthivorous	223	4	200000	rounded	flat	<sup>8</sup>	North Sea
<i>Mullus barbatus</i>	benthivorous	114	1.5	10000	forked	normal	<sup>23</sup>	Mediterranean
<i>Mullus surmuletus</i>	generalist	121	1.5*	10000**	forked	normal	<sup>23</sup>	Mediterranean
<i>Myoxocephalus quadricornis</i>	generalist	190	2**	10000***	truncated	normal	<sup>24</sup>	Barents Sea
<i>Myoxocephalus scorpius</i>	generalist	210	2	10000	rounded	normal	<sup>24</sup>	Barents Sea
<i>Myxine glutinosa</i>	benthivorous	293	3	100	continuous	eellike	<sup>25</sup>	Europe
<i>Pholis gunnellus</i>	benthivorous	176	2	100	rounded	eellike	<sup>1</sup>	Baltic Sea
<i>Platichthys flesus</i>	benthivorous	274	2	1000000	rounded	flat	<sup>11</sup>	Bay of Biscay
<i>Pleuronectes platessa</i>	benthivorous	259	2.5	552000	rounded	flat	<sup>26</sup>	Barents Sea
<i>Pollachius virens</i>	piscivorous	298	4.6	2900000	emarginated	normal	<sup>8</sup>	North Sea
<i>Salmo salar</i>	piscivorous	664	5	17000	truncated	normal	<sup>11</sup>	Europe
<i>Salmo trutta</i>	piscivorous	573	2.5	2000	truncated	normal	<sup>11</sup>	Europe
<i>Scophthalmus maximus</i>	generalist	296	3	5000000	rounded	flat	<sup>27</sup>	Baltic Sea
<i>Scophthalmus rhombus</i>	generalist	292	1.6	5000000	rounded	flat	<sup>28,29</sup>	Europe
<i>Solea solea</i>	benthivorous	240	3	300000	rounded	flat	<sup>30</sup>	North Sea
<i>Squalus acanthias</i>	piscivorous	693	30	6	emarginated	elongated	<sup>31</sup>	Baltic Sea
<i>Syngnathidae</i>	generalist	258	1	65****	rounded	eellike	<sup>32</sup>	Aegan Sea
<i>Trachinus draco</i>	generalist	215	1	57600	truncated	normal	<sup>33,34</sup>	Baltic Sea
<i>Trisopterus minutus</i>	generalist	145	1	10000	truncated	normal	<sup>35</sup>	Aegan Sea
<i>Zoarces viviparus</i>	benthivorous	227	2	100	continuous	eellike	<sup>36</sup>	Baltic Sea

\* As *Gobius niger* \*\* As *Mullus barbatus* \*\*\* As *Myoxocephalus Scorpius* \*\*\*\* As *Syngnathus acus*

Table S2. Initial traits database and the final 6 traits selected (bold).

Trophic level
<b>Trophic guilds (Diet )</b>
Maximum length (Lmax)
Length at 50% maturity (L50)
<b>Mean survey length (Lmean)</b>
<b>Age at 50% maturity (A50)</b>
Life expectancy
Offspring size
<b>Fecundity</b>
Schooling behaviour
<b>Caudal shape</b>
<b>Body shape</b>
Barbels presence

Table S3. Spatial autocorrelation of the residuals from the gam model: Mantel test (R package vegan)

Gam Model	Mantel statistic r	P-value
SR~ Salinity	0.0063	0.45
SR~ Salinity + habitat + Oxygen	0.0268	0.29
FR ~ Salinity	0.0063	0.39
FR ~ Salinity + habitat	0.0315	0.27

Table S4. Variables and parameters of the final generalized additive models (GAMs). Sign: whether the fitted relationships are predominately positive or negative; edf: estimated degrees of freedom for the model smooth terms (edf > 1 indicates a nonlinear relationship); DEV: degree of explained variance; Significance code: \*p<0.05, \*\*p<0.01, \*\*\* p<0.001.

Metrics	Best models and individual parameters	Sign	edf	DEV (%)
Species Richness	Best model: Salinity + Oxygen + habitat			92.5
	Salinity***	+	1.91	
	Oxygen***	+	0.94	
	Habitat***	-	0.94	
	Best single-variable model: Salinity			84.7
	Salinity***	+	1.88	
Functional Richness	Best model: Salinity + habitat			63.1
	Salinity***	+	1.72	
	Habitat***	-	0.87	
	Best single-variable model: Salinity			53.8
	Salinity***	+	1.81	
Functional Evenness	Best model: Depth sd + Depth + habitat			37.1
	Depth sd ***	+	1.8	
	Depth ***	+	1	
	Habitat *	□	0.7	
	Best single-variable model:			11.5
	Temperature *	□	1.2	
Functional Divergence	Best model: Depth sd			19.9
	Depth sd**	□	1.3	




Figure S1. No influence of the sampling effort on the species richness




Figure S2. Correlations and Pearson coefficient between the different length metrics (left) and age metrics (right)




Figure S3. Influence of the traits database chosen: correlation (Pearson) between functional richness calculated with the 13 traits compared to functional richness calculated with the 6 selected traits only.




Figure S4. Species Richness and the three facets of functional diversity<sup>37</sup> of the Baltic Sea demersal fish communities. The values are cut in quintiles.




Figure S5. Temporal study of the relationship between SR and FRic. The years from 2004 to 2012 are represented using a grey scale (from dark in 2004 to light grey in 2012).




Figure S6. Rao Q versus wFDiss. Rao Q<sup>38</sup> represents the average functional dissimilarity, i.e. distance, of two individuals randomly selected in a community, while wFDiss represents the average functional dissimilarity of two individuals from two different species in a community. Hence, Rao Q is always lower than wFDiss as it takes into account the functional dissimilarity of two individuals from the same species ( $d_{ii}=0$ ). Green colors represent lower Rao Q or wFDiss values, whilst red colors represents higher values. The black points represents the single hauls from which a smoothing function was applied (thin plate spline regression from fields package in R).

## References

1. Muus, B. J. & Nielsen, J. G. Sea fish. Scandinavian Fishing Year Book, Hedehusene, Denmark. 340 p. (1999).
2. Walker, P. Sensitive skates or resilient rays? Spatial and temporal shifts in ray species composition in the central and north-western North Sea between 1930 and the present day. *ICES Journal of Marine Science* **55**, 392–402 (1998).
3. Gauld, J. A. & Hutcheon, J. R. Spawning and fecundity in the lesser sandeel, *Ammodytes marinus* Raitt, in the north-western North Sea. *Journal of Fish Biology*, 36: 611–613. (1990).
4. MacNamara, R. & McCarthy, T. K. Size-related variation in fecundity of European eel (*Anguilla anguilla*). *ICES Journal of Marine Science* **69**, 1205–1217 (2012).
5. Vollestad, V. Geographic variation in age and length at metamorphosis of maturing European eel: environmental effects and phenotypic plasticity. *Journal of Animal Ecology* **61**, 41–48 (1992).
6. Gibson, R. N. & Ezzi, I. a. The biology of the sculdfish, *Arnoglossus laterna* (Walbaum) on the west coast of Scotland. *Journal of Fish Biology* 575 (1980). doi:10.1111/j.1095-8649.1980.tb02788.x
7. Déniel, C. La reproduction des poissons plats (téléostéens-pleuronectiformes) en baie de Douarnenez. 2. Cycles sexuels et fécondité des Soleidae: *Solea vulgaris vulgaris*, *Solea lascaris*, *Buglossidium luteum* et *Microchirus variegatus* = Flatfish reproduction in Douarnen. *Cah. Biol. Mar.* **25(3)**, 257–285 (1984).
8. Jennings, S., Greenstreet, S. P. R. & Reynolds, J. D. Structural change in an exploited fish community: A consequence of differential fishing effects on species with contrasting life histories. *Journal of Animal Ecology* **68**, 617–627 (1999).
9. Işmen, A., Işmen, P. & Başusta, N. Age, growth and reproduction of tub gurnard (*Chelidonichthys lucerna* L. 1758) in the Bay of İskenderun in the Eastern Mediterranean. *Turkish Journal of Veterinary and Animal Sciences* **28**, 289–295 (2004).
10. Davenport, J. Synopsis of biological data on the lumpsucker, *Cyclopterus lumpus* (Linnaeus, 1758). *Food & Agriculture Org.* **147**, (1985).
11. Froese, R. & Pauly, D. (Eds.), Fishbase. World Wide Web Electronic Publication, 2012., [www.fishbase.org](http://www.fishbase.org) (2012). at <[www.fishbase.org](http://www.fishbase.org)>
12. Kosior, M., Trella, K. & Jaworski, A. Fecundity of cod in the Southern Baltic in the late 1990s. *Acta Ichthyol. Piscat* **31**, 55–75 (2001).
13. Burnett, J., Ross, M. R. & Clark, S. H. Several biological aspects of the witch flounder (*Glyptocephalus cynoglossus* (L.)) in the Gulf of Maine-Georges Bank region. *Journal of Northwest Atlantic Fishery Science* **12**, 15–25 (1992).
14. Fossen, I., Albert, O. T. & Nilssen, E. M. Back-calculated individual growth of long rough dab (*Hippoglossoides platessoides*) in the Barents Sea. *ICES Journal of Marine Science* 689–696 (1999). doi:10.1006/jmsc.1999.0486
15. Robson, S. Age, growth, reproductive biology and population dynamics of the common megrim *Lepidorhombus whiffiagottis* (Walbaum, 1792) from off the west coast of Ireland. *Unpublished thesis (Doctor of Philosophy in Fisheries Biology), Galway-Mayo Institute of Technology* (2004).
16. Rijnsdorp, a. D., Vethaak, a. D. & Van Leeuwen, P. I. Population biology of dab *Limanda limanda* in the southeastern North Sea. *Marine Ecology Progress Series* **91**, 19–35 (1992).
17. Wienerroither, R. Atlas of the Barents Sea fishes. *IMR/PINRO Joint Report* **1**, 1–272. (2011).
18. Marty, L., Rochet, M. J. & Ernande, B. Temporal trends in age and size at maturation of four North Sea gadid species: Cod, haddock, whiting and Norway pout. *Marine Ecology Progress Series* **497**, 179–197 (2014).
19. Trippel, E. a & Neil, S. R. Maternal and seasonal differences in egg sizes and spawning activity of northwest Atlantic haddock (*Melanogrammus aeglefinus*) in relation to body size and condition. *Canadian Journal of Fisheries and Aquatic Sciences* **61**, 2097–2110 (2004).
20. Gerritsen, H. D., Armstrong, M. J., Allen, M., McCurdy, W. J. & Peel, J. a D. Variability in maturity and growth in a heavily exploited stock: whiting (*Merlangius merlangus* L.) in the Irish Sea. *ICES Journal of Marine Science* **61**, 98–112 (2004).

21. Pineiro, C. & Sainza, M. Age estimation, growth and maturity of the European hake (*Merluccius merluccius* (Linnaeus, 1758)) from Iberian Atlantic waters. *ICES Journal of Marine Science: Journal du Conseil* **3139**, 1352–1360 (2003).
22. Murua, H., Lucio, P., Santurtún, M. & Motos, L. Seasonal variation in egg production and batch fecundity of European hake *Merluccius merluccius* (L.) in the Bay of Biscay. *Journal of Fish Biology* **69**, 1304–1316 (2006).
23. Tirasin, E., Unluoglu, A. & Cihangir, B. Fecundity of red mullet (*Mullus barbatus* L., 1758) along the Turkish coasts of the Mediterranean sea. (2007).
24. Luksenburg, J. a., Pedersen, T. & Falk-Petersen, I. B. Reproduction of the shorthorn sculpin *Myoxocephalus scorpius* in northern Norway. *Journal of Sea Research* **51**, 157–166 (2004).
25. Powell, M. L., Kavanaugh, S. I. & Sower, S. a. Current knowledge of hagfish reproduction: implications for fisheries management. *Integrative and comparative biology* **45**, 158–165 (2005).
26. Christiansen, J. S. Maternal output in polar fish reproduction. *Fishes of Antarctica*. Springer Milan 41-52. 1998
27. Nissling, A., Florin, A.-B., Thorsen, A. & Bergström, U. Egg production of turbot, *Scophthalmus maximus*, in the Baltic Sea. *Journal of Sea Research* **84**, 77–86 (2012).
28. Van der Hammen, T. *et al.* Population ecology of turbot and brill: What can we learn from two rare flatfish species? *Journal of Sea Research* **84**, 96–108 (2013).
29. Félix, P. M., Vinagre, C. & Cabral, H. N. Life-history traits of flatfish in the Northeast Atlantic and Mediterranean Sea. *Journal of Applied Ichthyology* **27**, 100–111 (2011).
30. Witthames, P. R. & Walker, M. G. Determinacy of fecundity and oocyte atresia in sole (*Solea solea*) from the Channel, the North Sea and the Irish Sea. *Aquatic Living Resources* **8**, 91–109 (1995).
31. Stenberg, C. Life History of the Piked Dogfish (*Squalus acanthias* L.) in Swedish Waters. *J. Northw. Atl. Fish. Sci.*, 35: 155-164. (2005).
32. Gurkan, S., Taskavak, E. & Hossucu, B. The reproductive biology of the great pipefish *Syngnathus acus* (Family: Syngnathidae) in the Aegean Sea. *North-Western Journal of Zoology* **5**, 179–190 (2009).
33. Ak, O. & Genç, Y. Growth and reproduction of the greater weever ( *Trachinus draco* L ., 1758 ) along the eastern coast of the Black Sea. **19**, 95–110 (2013).
34. Bagge, O. The biology of the greater weever (*Trachinus draco*) in the commercial fishery of the Kattegat. *ICES Journal of Marine Science* **61**, 933–943 (2004).
35. Metin, G., Ilkyaz, A. T. & Kinacigil, H. T. Growth, mortality, and reproduction of poor cod (*Trisopterus minutus* Linn., 1758) in the central Aegean sea. *Turkish Journal of Zoology* **32**, 43–51 (2008).
36. Strand, J., Andersen, L., Dahllöf, I. & Korsgaard, B. Impaired larval development in broods of eelpout (*Zoarces viviparus*) in Danish coastal waters. *Fish Physiology and Biochemistry* **30**, 37–46 (2004).
37. Villéger, S., Mason, N. W. H. & Mouillot, D. New multidimensional functional diversity indices for a multifaceted framework in functional ecology. *Ecology* **89**, 2290–301 (2008).
38. Rao, C. R. Diversity and dissimilarity coefficients: A unified approach. *Theoretical Population Biology* **21**, 24–43 (1982).